

Intensive harvesting menaces trees producing fodder, edible fruit, and gum in Abu Gadaf natural reserved forest, Sudan

Elmugheira M. I. Mohammed¹, Elmalih M. I. Mohammed² and Bashir A. A. Gerbeel³

¹Department of Forest Management Science, Faculty of Forest Sciences and Technology, University of Gezira, Wad Medani, Sudan

²Freelance, Gedaref State, Sudan

³Forests National Corporation, Ministry of Agriculture and Forests, Blue Nile Region, Sudan

ABSTRACT

Background: This study aimed to assess the influence of intensive harvesting on tree dendrometric parameters, seedling recruitment, and sapling density of identified tree species in Abu Gadaf natural reserved forest, Sudan.

Methods: We used a stratified sampling design for data collection, and the identified tree species were classified into five groups based on the main product utilized locally. Data were collected in 46 sample plots of 1000m² systematically distributed across the upper and lower land sites of Abu Gadaf natural reserved forest. Analysis of variance, paired-samples t-test, and regressions in JAMovi and Minitab were performed for data analysis.

Results: Findings revealed that tree-producing gum and edible fruits have small diameters compared to fodder and building pole-producing ones. Although *Acacia polyacantha* and *Acacia seyal* have larger mean diameter in the lowerland sites, *Sterculia africana* and *Sterculia setigera* displayed the inverse pattern with significant differences between sites ($F_{1,45}=102.7$ and $p<0.01$; $F_{1,45}=108.3$ and $p=0.01$; $F_{1,45}=120.2$ and $p=0.03$, $F_{1,45}=111.2$ and $p=0.02$, respectively). Moreover, the mean volume of *Anogeissus leiocarpus*, *Balanites aegyptiaca*, *Terminalia brownii*, and *Sclerocarya birrea* in upperland sites was twice, three, and five times equal to that of lowerland ones, respectively.

Conclusion: We concluded that intensive harvesting severely affected fodder, edible fruit, and gum-producing trees in the reserve and reduced their populations. We further recommend an introduction of the Taungya system concept for the restoration of the degraded sites.

KEYWORDS

Agroforestry; Biosphere reserve; Firewood; Forest conservation; Natural regeneration

ARTICLE HISTORY

Received 25 August 2023;

Revised 18 October 2023;

Accepted 25 October 2023

Introduction

Trees play significant ecological and socio-economic roles through soil enrichment, food provision, income generation, and carbon sequestration [1,2]. All tree parts, including fruits, leaves, twigs, branches, stems, and roots, are frequently utilized to satisfy innumerable needs and services [3,4]. Tree crown properties such as crown size and shape, leaves phyllotaxy, twigs density, and branches arrangement with other edaphic and growth factors determine tree growth activities [5], fruits and seeds production [6]. They also govern the shading area and shade quality [7], reduction of air pollutants, rain filtering, and running-water speed [8,9], wind-breaking efficiency [10,11], and carbon sink adequacy [12]. Moreover, the total tree height, diameter at breast height, and seedlings density are core components for tree volume calculation [13,14], site quality index [12,15], and regeneration efficiency [16,17], respectively. However, over-utilization can interrupt these functions and limit their roles.

Gum-producing trees like *Acacia polyacantha*, *Acacia seyal*, *Acacia senegal*, *Boswellia papyrifera*, and *Sterculia setigera* received a considerable number of studies related to seedling propagation, tapping techniques, ecological range, and their potential to enhance soil fertility [18-21]. Yet, limited literature is available on how intensive harvesting and over-utilization can

affect their population dynamics and sustainability. Similar scenarios are observed for fodder, fruit, and pole-producing trees. The study conducted by Dampney et al. concluded that high fodder production in natural forests is strongly associated with other biomass proxies and beneficial arthropods [22]. However, carbon storage potential, population structure, and stand density can dramatically decline due to human stresses and extreme environmental conditions [13,22-24]. Moreover, Benítez-Badillo et al. documented that land use changes influenced the distribution of wild edible fruits and deteriorated the tropical forests of Veracruz State in Mexico [25]. Therefore, studies dealing with the influence of over-harvesting, shifting cultivation, overgrazing, and mining activities on forest trees contribute significantly to the sustainable management of forest resources [22,26-28].

Over-harvesting trees and other forest products reduce forest biodiversity and accelerate land degradation in many continents, including Africa [29-31]. In the Sahel area, with a high percentage of livestock keeping and agricultural activities, reduction in vegetation cover, soil fertility, species richness, and forest yield is typical [2,32-34]. Such reduction can threaten the livelihoods of rural communities in the Sahel,

*Correspondence: Elmugheira M. I. Mohammed, Department of Forest Management Science, Faculty of Forest Sciences and Technology, University of Gezira, Wad Medani, Sudan, e-mail: elmugheira1984@gmail.com

© 2023 The Author(s). Published by Reseapro Journals. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

particularly the vulnerable societies in Sudan that principally depend on natural forests. One of these forests is Abu Gadaf natural reserved forest.

Abu Gadaf is a natural reserve found in the Blue Nile region, offering numerous functions and services to more than five thousand families living in four villages surrounding the forest [35-37]. The locals within and around this forest harvest timbers for firewood and charcoal production [38,39], green twigs and leaves for livestock feeding [40,41], bark and roots for traditional medicine [35,42], and fruits for food and income generation [36,43]. While harvesting activities are dramatically increasing, information on how this utilization can affect the tree dendrometric parameters, seedlings recruitment, and saplings development is limited. Aiming to fill this gap, we assessed the influence of intensive harvesting on crown properties, tree height, and seedlings density of identified tree species in Abu Gadaf natural reserved forest, Sudan.

The study compared the tree diameter at breast height (DBH), total tree height, crown width, and tree volume between the upperland sites (Mountainous lands) and lowerland sites (Flatlands). We further assessed the seedling, sapling, and adult tree densities within and across the study sites. Additionally, we set two hypotheses. Firstly, lowerland sites have limited seedling and sapling densities with small crown width in comparison to upperland sites. Secondly, the upperland sites have larger tree DBH, total tree height, and tree volume compared to lowerland ones.

Materials and Methods

Study area

The Abu Gadaf Natural Reserved Forest is located at 11° 25' 00" N, 11° 31' 00" N, 34° 50' 00" E, and 34° 55' 00" E (Figure 1), covering an area of 4,413.87 ha [35,44]. The forest hosts useful food and feed tree species such as *Acacia senegal*, *Anogeissus leiocarpus*, *Balanites aegyptiaca*, *Boswellia papyrifera*, *Combretum hartmannianum*, *Grewia mollis*, *Lannea fruticosa*, *Tamarindus indica*, *Terminalia brownii*, and *Ziziphus spina-christi* [35,43]. The average monthly rainfall differs from <50mm in winter to >300mm in autumn, with a minimum and maximum temperature of 23 °C and 44 °C, respectively [35,36,43]. Moreover, the reserve topography and soil types are sandy soil in the upperland sites and clay soil in the lowerland sites [13,45]. During the rainy season, some wild animals like *Cercopithecus aethiops*, *Numida meliagris*, *Phacochoerus aethiopicus*, *Redunca redunca*, and *Struthio camelus* move from Dinder Biosphere Reserve to Abu Gadaf forest [45,46].

Data collection

We used the stratified sampling design for data collection. The study area was divided based on the forest topography into upper and lower land sites, where 46 sample plots of 1000m² were systematically established across the sites. After determining the plot boundary and identifying the tree species, tree diameter (diameter at breast height), total tree height, bole height, and crown height, were measured using diameter tape, Suunto clinometer, and Spiegel Relaskop, respectively [47,48]. The identified tree species were classified into five groups as gum-producing trees, edible fruit-producing trees, building pole-producing trees,

fodder-producing trees, and firewood-producing trees. The classification is based on the main tree product utilized by the local community. The crown diameter was measured as described by Mohammed et al. and Hassan et al. [13,45]. All cruised woody plants (tree species) were distinguished into seedlings (<3cm in stem diameter), saplings (≥ 3cm to <7cm in stem diameter), and adult trees (≥ 7cm in stem diameter) as documented by Mohammed et al., Gebeyehu et al. and Ligate et al. [46,49,50].




Figure 1. The map of Abu Gadaf Natural Reserved Forest and its stratified upper and lowerland sites.

Data analysis

We checked for normality using the Shapiro test and calculated the tree density and volume using equations 1 and 2 [13,47]. We further compared the crown width, height, and diameter at breast height values of different tree species within the sites and between the upper and lower land sites using a two-way ANOVA in JAMovi [43,48]. The paired-sample t-test was performed in IBM SPSS to compare the density of seedlings, saplings, and mature trees [14,51]. Tukey post hoc test with a p-value of 0.05 was applied for significant differences as recommended by Gebeyehu et al., Maua et al. and Hasoba et al. [12,17,52]. Moreover, correlations between seedlings density and the No. of tree stumps at the upper and lowerland sites were run in Minitab 17 [53-55].

Tree density= Number of tree stems/area sampled----- (1)

Tree volume=Tree basal area × total tree height × form factor----- (2)

Results

Tree diameter and height

The trees producing gum and edible fruits have the small diameter at breast height compared to fodder and building pole-producing ones (Figure 2). While *Acacia polyacantha* and *Acacia seyal* illustrated larger mean diameter in the lowerland sites, *Sterculia africana* and *Sterculia setigera* displayed the inverse pattern with significant differences between sites ($F_{1,45}=102.7$ and $p<0.01$; $F_{1,45}=108.3$ and $p=0.01$; $F_{1,45}=120.2$ and

$p=0.03$, $F_{1,45}=111.2$ and $p=0.02$, respectively, Figure 2). Moreover, the mean total height of *Anogeissus leiocarpus*, *Combretum hartmannianum*, *Terminalia brownii*, and *Ziziphus spina-christi* in the upperland site was double that of lowerland site with significant differences across sites ($F_{1,45}=97.2$ and $p<0.01$; $F_{1,45}=101.3$ and $p=0.01$; $F_{1,45}=89.8$ and $p=0.01$, $F_{1,45}$

$=133.2$ and $p=0.01$, respectively, Figure 3). However, *Acacia senegal* and *Maerua angolensis* showed no significant differences between upper and lowerland sites for both diameter and total tree height (Figures 2 and 3).

Tree crown width and volume




Figure 2. Mean diameter at breast height (cm) for the identified five groups of tree species, classified based on the main utilized product. (A) Gum-producing trees, (B) Edible fruit-producing trees, (C) Building pole-producing trees, (D) Fodder-producing trees, and (E) Firewood-producing trees. Asterisks on the bar illustrate the significant differences between sites for various tree species based on Tukey post hoc test ($\alpha=0.05$) where $*=p<0.05$ and $**=p<0.01$.




Figure 3. Mean total tree height (m) for the identified five groups of tree species, classified based on the main utilized product. (A) Gum-producing trees, (B) Edible fruit-producing trees, (C) Building pole-producing trees, (D) Fodder-producing trees, and (E) Fire-producing trees. Asterisks on the bar illustrate the significant differences between sites for various tree species based on Tukey post hoc test ($\alpha=0.05$) where $*=p<0.05$ and $**=p<0.01$.

The mean crown width of *Acacia seyal*, *Combretum ghasalense*, and *Combretum glutinosum* in the upperland site were three times equal to that of the lowerland site and significantly differed between sites ($F_{1, 45}=82.3$ and $p<0.001$; $F_{1, 45}=128.6$ and $p=0.01$; $F_{1, 45}=113.7$ and $p=0.01$, respectively, Figure 4). Though all fodder and firewood-producing trees showed significant differences between sites, *Boswellia papyrifera*, *Combretum micranthum*, *Combretum molle*, *Dalbergia melanoxylon*, *Entada africana*, and *Sterculia africana* exhibited no significant differences across the sites (Figure 4). However, *Acacia senegal* in comparison to *Adansonia digitata* and *Tamarindus indica* displayed an inverse trend with large tree crown in lowerland one ($F_{1, 45}=142.3$ and $p<0.04$, Figure 4).

Furthermore, the mean volume of *Anogeissus leiocarpus*, *Balanites aegyptiaca*, *Terminalia brownii*, and *Sclerocarya birrea* in upperland site were two, three, and five times equal to that of lowerland one, respectively ($F_{1, 45}=91.4$ and $p<0.01$; $F_{1, 45}=78.6$ and $p=0.001$; $F_{1, 45}=73.8$ and $p=0.001$, $F_{1, 45}=81.1$ and $p=0.001$, respectively, Figure 5). In addition to that the average

volume of *Lannea kerstignii* and *Lannea nigritana* in the upperland site, were more than six times to that of lowerland site with significant differences between sites ($F_{1, 45}=119.2$ and $p<0.001$; $F_{1, 45}=116.1$ and $p=0.001$, respectively, Figure 5). While *Acacia polyacantha* and *Grewia flavescens* have the minimum mean volume ($<0.09\text{m}^3$), *Adansonia digitata* and *Sterculia setigera* attained the maximum mean volume for both lower and upperland sites ($>1.4\text{m}^3$ and 2.3m^3 , respectively, Figure 5).

Based on the group classification, trees producing building poles and firewood had lower mean volume compared to the other three groups (Figure 5). The average volume of *Acacia senegal* in the upperland site was half of lowerland one with significant differences between sites ($F_{1, 45}=79.7$ and $p<0.03$, Figure 5). However, *Pterocarpus lucens*, *Stereospermum kunthianum*, *Terminalia macroptera*, and *Ziziphus spina-christi* demonstrated an opposite scenario with higher values of mean volume in the upperland site compared to another one (Figure 5).




Figure 4. Mean crown width (m) for the identified five groups of tree species, classified based on the main utilized product. (A) Gum-producing trees, (B) Edible fruit-producing trees, (C) Building pole-producing trees, (D) Fodder-producing trees, and (E) Firewood-producing trees. Asterisks on the bar illustrate the significant differences between sites for various tree species based on Tukey post hoc test ($\alpha = 0.05$) where *= $p < 0.05$, **= $p < 0.01$, and ***= $p < 0.001$.




Figure 5. Mean tree volume (m³) for the identified five groups of tree species, classified based on the main utilized product. (A) Gum-producing trees, (B) Edible fruit-producing trees, (C) Building pole-producing trees, (D) Fodder-producing trees, and (E) Firewood-producing trees. Asterisks on the bar illustrate the significant differences between sites for various tree species based on Tukey post hoc test ($\alpha = 0.05$) where *= $p < 0.05$, **= $p < 0.01$, and ***= $p < 0.001$.

Seedlings, saplings, and adult trees density and volume

The correlation between the no. of tree stumps and mean seedlings density of gum-producing, edible fruit-producing, building pole-producing, fodder-producing, and firewood-producing trees in the upperland site demonstrated strong positive relationships with $R^2 = 0.99$ ($\beta = 6.7$ and $p = 0.03$; $\beta = 5.4$ and $p = 0.04$; $\beta = 7.4$ and $p = 0.01$; $\beta = 4.7$ and $p = 0.04$; $\beta = 7.2$ and $p = 0.01$; respectively, Figure 6). However, for lowerland site,

it formed an inverse pattern with $R^2 = 0.97$ ($\beta = 1.9$ and $p = 0.01$; $\beta = 6.8$ and $p = 0.03$; $\beta = 1.3$ and $p = 0.01$; $\beta = 11.2$ and $p = 0.04$; $\beta = 2.6$ and $P = 0.01$; respectively, Figure 6). Among the five groups, fodder and firewood-producing trees have the lowest and highest seedlings and saplings densities, respectively (Table 1). The saplings density of firewood-producing trees in the upperland site was two times that of edible fruits, fodder, and firewood producing trees at lowerland site (Table 1). Similar trends were observed for seedlings, saplings, and mature trees volume (Table 1).




Figure 6. Relationship between the no. of tree stumps (stump ha⁻¹) and seedlings density for gum-producing trees, edible fruit-producing trees, building material-producing trees, fodder-producing trees, and firewood-producing trees in (A) Upperland site and (B) Lowerland site of Abu Gadaf natural reserved forest, Sudan.

Table 1. Mean density ($\pm$ SD) and volume ($\pm$ SD) for the identified seedlings, saplings, and mature trees of the five classified groups of tree species inventoried in Abu Gadaf natural reserved forest, Sudan.

Group name	Seedlings			Saplings			Mature trees		
	Upperland	Lowerland	P	Upperland	Lowerland	P	Upperland	Lowerland	P
Density (stem/ha)									
Gum-producing trees	76 $\pm$ 0.4	58 $\pm$ 2.6	0.04	72 $\pm$ 0.3	51 $\pm$ 1.4	0.00	64 $\pm$ 0.8	42 $\pm$ 2.7	0.03
Edible fruit-producing trees	74 $\pm$ 0.6	57 $\pm$ 1.9	0.03	63 $\pm$ 4.8	41 $\pm$ 0.6	0.02	45 $\pm$ 1.6	22 $\pm$ 0.6	0.00

Building pole-producing trees	82 ± 3.5	60 ± 0.9	0.00 1	74 ± 0.5	53 ± 0.7	0.00 1	52 ± 2.8	33 ± 0.7	0.02
Fodder-producing trees	66 ± 1.3	50 ± 0.7	0.03	65 ± 1.5	43 ± 0.2	0.00 1	43 ± 0.9	21 ± 1.2	0.00 1
Firewood-producing trees	85 ± 3.4	58 ± 1.8	0.00 1	85 ± 5.4	42 ± 0.5	0.00 1	65 ± 2.9	42 ± 0.9	0.03
Volume (m ³)									
Gum-producing trees	0.811 ± 0.02	0.521 ± 0.01	0.02	0.752 ± 0.02	0.442 ± 0.01	0.01	0.655 ± 0.02	0.546 ± 0.03	0.03
Edible fruit-producing trees	0.789 ± 0.01	0.543 ± 0.01	0.03	0.951 ± 0.02	0.531 ± 0.02	0.02	0.850 ± 0.02	0.562 ± 0.03	0.02
Building pole-producing trees	0.565 ± 0.02	0.311 ± 0.02	0.02	0.498 ± 0.01	0.256 ± 0.02	0.01	0.479 ± 0.03	0.298 ± 0.02	0.01
Fodder-producing trees	0.624 ± 0.01	0.351 ± 0.01	0.01	0.587 ± 0.02	0.321 ± 0.01	0.03	0.543 ± 0.02	0.224 ± 0.03	0.02
Firewood-producing trees	0.721 ± 0.03	0.335 ± 0.02	0.01	0.643 ± 0.01	0.312 ± 0.03	0.01	0.492 ± 0.01	0.215 ± 0.02	0.01

Discussion

Tree diameter and height

We found that trees locally used for the production of building poles and firewood have large DBH in comparison to gum, edible fruits, and fodder-producing trees. Nevertheless, the lowerland site host limited adult and juvenile trees than the upperland one. Similar results were reported by Beche et al., Hassan, Adam et al. and Ereso [20,36,56,57]. The leading species of gum-producing trees are *Acacia senegal*, *Acacia seyal*, and *Boswellia papyrifera*, while for edible-fruit trees, *Balanites aegyptiaca*, *Grewia bicolor*, *Grewia flavescens*, and *Grewia mollis* are on top. Though *Grewia* sp. are generally shrubs to small trees, *Balanites aegyptiaca* and *Boswellia papyrifera* are sensitive to intensive debarking and debranching, respectively [20,58,59]. Moreover, selective logging of codominant and free-growing trees can eliminate the trees with large diameter and fruit-producing trees [1,27,60]. Researchers, concluded that over-harvesting of mature and dominant trees in the forest and biosphere reserves disturbed the forest stand composition in Ethiopia, Niger, Tanzania, and Kenya, respectively [17,50,61,62].

The findings of total tree height analysis shown that mean height at the lowerland site was half of that in the upperland site for most cruised tree species in Abu Gadaf natural reserved forest [24,28,35]. While Ji et al., Chaudhary et al. and Adekunle and Olagoke referred the low tree height to unauthorized timber harvesting [63-65], Maua et al., Mohammed et al. and Fakhry and Aljedaani related it to both intensive livestock browsing and illegal logging [17,46,66]. The mountainous terrain of the upperland site and its farness from the locals' residences increased the number of mature trees, thus the tree diameter and height. The rough terrain, site accessibility, and utilization purpose can enhance or worsen the species conservation status and population structure [67-69]. However, the current reduction in tree diameter and total height at the lowerland site

of Abu Gadaf elucidates the severe effects of illegal logging and debranching applied by the forest- nearby communities that call for quick intervention and restoration programs. Vulnerable tree species like *Grewia bicolor*, *Grewia flavescens* and *Maerua angolensis* need urgent conservation action and detailed study on their spatial distribution and juvenile growth rate.

Though literature documented that, anthropogenic activities like livestock keeping, farming, and charcoal production can negatively affect forest trees [66,70-72], this is the first time to have specific information on gum, fodder, and edible fruit-producing trees, particularly for natural Sudanese forests.

Tree crown width and volume

The study findings revealed that the lowerland site accommodates trees with small crown width and volume compared to the upperland site. The small crown width in Abu Gadaf forest, particularly for fodder-producing trees, can be referred to over-browsing by livestock and frequent removal of fresh twigs and branches for sheep and goat feeding. Studies performed by Hassan et al., Fakhry and Aljedaani and Bondé et al. stated that camel browsing reduced the crown width of fodder trees at highly disturbed sites in Burkina Faso, Saudi Arabia, and Sudan to less than 40%, 60%, and 35%, respectively [45,66,73]. Moreover, Mohammed et al., Ahmed and Desougi, Hassan et al. and Ali et al. reported that illegal cutting of forest trees deteriorated the population of *Acacia senegal*, *Acacia seyal*, *Combretum hartmannianum*, and *Terminalia brownii* in Tozi, Hegleig, Okalma, and Dinder reserves in Sudan, respectively [13,18,45,74].

Furthermore, the decline in tree volume is directly associated with the reduction in tree height and diameter at breast height which are strongly linked to the site quality, biotic and abiotic disturbances, and other environmental conditions [11,24,33,75]. Intensive harvesting of mother and old-growing

trees results in limited upper story trees [5,76], lower importance value index [2,15,77], smaller basal area [78,79], and consequently tree volume [13]. Therefore, the limited volume of firewood and pole-producing trees in Abu Gadaf natural reserves forest originates from the current observed illegal logging activity and other anthropogenic deeds like livestock keeping, agriculture, and urbanization. Likewise, outcomes were concluded by Crump et al., Blackham et al. and Ewunetu et al. [80-82].

Seedlings, saplings, and adult trees density and volume

The efficient natural regeneration of forest trees depends on seed viability, suitable edaphic and climatic conditions for seed germination, site competition, and biotic and abiotic disturbances [14,83,84]. Stands with high anthropogenic disturbances, poor soil fertility, and low precipitation rate attain poor seed germination rate, lower seedling growth rate, and consequently limited sapling recruitment [85-87]. Our findings support the previously documented results and highlight the significant influence of human stresses on seedling density and saplings recruitment. The strong positive relationship between the number of tree stumps and seedlings density in the upperland of Abu Gadaf natural reserved forest illustrated that this site is subject to less crown debranching, thus more seed production resulting in more seedlings in the newly opened space. However, for the inverse pattern at the lowerland site, the intensive removal of fruiting branches reduced the number of produced seeds, consequently, seedling and sapling density. Various studies are aligned with these findings [51,88,89].

While browsing commonly affects adult and upper story trees, illegal harvesting impacts mature trees and regeneration [90-92]. This study is among the few that elucidated how intensive harvesting and frequent logging of adult trees can influence the density of seedlings, saplings, and adult trees. The lower density of fodder and edible fruit-producing trees in Abu Gadaf is directly associated with the intensive utilization of these species for food, feed, medicine, and income generation, resulting in a lower density of adult and juvenile trees and demand for a conservation policy and plan. Studies conducted by Aubad et al., Kutnar et al., Tetemke et al. and Wang et al. indicated that combined effects of livestock grazing and logging reduced the species richness, stocking density of adult trees, regeneration ratio, and population structure [93-96]. The species like *Adansonia digitata*, *Balanites aegyptiaca*, *Hyphaena thebaica*, *Tamarindus indica*, and *Ziziphus spina-christi* need to be promoted towards the enhancement of local community income and moderation of the increasing pressure on other tree species at the reserve.

Conclusions

We concluded that intensive harvesting by local communities within and around Abu Gadaf natural reserved forest had decreased the population of woody plants, particularly trees producing fodder, edible fruits, and gum. The lowerland site hosts trees with small diameters, crown widths, heights, and volumes compared to the upperland site. Even though *Grewia bicolor* and *Grewia flavescens* demand urgent intervention and conservation plan, *Adansonia digitata*, *Balanites aegyptiaca*, and *Ziziphus spina-christi* exhibited a potential source for future afforestation programs and locals livelihoods enhancement. However, more patrolling monitoring in the

lowerland site is needed to control the observed human interference and paves the road for the natural restoration of the degraded area. Additionally, an introduction of the Taungya system concept can guide the afforestation programs and rehabilitates the affected sites.

Acknowledgements

The authors thank the officers of the Forests National Corporation in the Blue Nile region for their kind support and genuine facilities during the fieldwork activities and data sorting. Special thanks to Prof. Talaat D. Abdel Magid, Prof. Hisham M. Tahir, Dr. Abass M. Elhag, and Dr. Yousif Suleiman for their guidance, support, and encouragement. We are also grateful to University of Gezira, Faculty of Forest Sciences and Technology, for the study permit.

Disclosure statement

No potential conflict of interest was reported by the authors.

References

1. Lindenmayer DB, Laurance WF. The ecology, distribution, conservation and management of large old trees. *Biol Rev.* 2017;92:1434-1458.
2. Neyia T, Neyia O, Abunyewa AA. Agroforestry parkland profiles in three climatic zones of Burkina Faso. *Int J Biol Chem Sci.* 2019;12(5):2119-2131.
3. Abdelaziz SM, Lemine FM, Tfeil HO, Filali-Maltouf A, Boukhary AO. Phytochemicals, antioxidant activity and ethnobotanical uses of *balanites aegyptiaca* (L.) del. fruits from the arid zone of mauritania, Northwest Africa. *Plants.* 2020;9(3):401.
4. Simons NK, Felipe-Lucia MR, Schall P, Ammer C, Bauhus J, Blüthgen N, et al. National forest inventories capture the multifunctionality of managed forests in Germany. *For Ecosyst.* 2021;8(1):1-9.
5. Di Salvatore U, Marchi M, Cantiani P. Single-tree crown shape and crown volume models for *Pinus nigra* J. F. Arnold in central Italy. *Ann For Sci.* 2021;78:1-10.
6. Mohammed EM, Hamed AM, Ndakidemi PA, Treydte AC. Illegal harvesting threatens fruit production and seedling recruitment of *Balanites aegyptiaca* in Dinder Biosphere Reserve, Sudan. *Glob Ecol Conserv.* 2021;29:e01732.
7. Hammond ME, Pokorný R, Okae-Anti D, Gyedu A, Obeng IO. The composition and diversity of natural regeneration of tree species in gaps under different intensities of forest disturbance. *J For Res.* 2021;32:1843-1853.
8. Derroire G, Tigabu M, Odén PC, Healey JR. The Effects of Established Trees on Woody Regeneration during Secondary Succession in Tropical Dry Forests. *Biotropica.* 2016;48(3):290-300.
9. Hofhansl F, Chacón-Madriral E, Fuchslueger L, Jenking D, Morera-Beita A, Plutzer C, et al. Climatic and edaphic controls over tropical forest diversity and vegetation carbon storage. *Sci Rep.* 2020;10(1):5066.
10. Thom D, Seidl R. Natural disturbance impacts on ecosystem services and biodiversity in temperate and boreal forests. *Biol Rev.* 2016;91(3):760-781.
11. Erdenebileg E, Wang C, Ye X, Cui Q, Du J, Huang Z, et al. Multiple abiotic and biotic drivers of long-term wood decomposition within and among species in the semi-arid inland dunes: A dual role for stem diameter. *Funct Ecol.* 2020;34(7):1472-1484.
12. Gebeyehu G, Soromessa T, Bekele T, Teketay D. Carbon stocks and factors affecting their storage in dry Afromontane forests of Awi Zone, northwestern Ethiopia. *J Ecol Environ.* 2019;43(1):1-18.
13. Mohammed EM, Hamid EA, Ndakidemi PA, Treydte AC. The stocking density and regeneration status of *Balanites aegyptiaca* in Dinder Biosphere Reserve, Sudan. *Trees For People.* 2022;8:100259.
14. Kingazi N, Petro R, Mbawambo JR, Munishi LK. Performance of *Pinus patula* in areas invaded by *Acacia mearnsii* in Sao Hill forest

- plantation, Southern Tanzania. *J Sustain For.* 2021;40(5):518-527.
15. Assogbadjo AE, Kakaï RL, Sinsin B, Pelz D. Structure of *Anogeissus leiocarpa* Guill., Perr. natural stands in relation to anthropogenic pressure within Wari-Marô Forest Reserve in Benin. *Afr J Ecol.* 2010;48(3):644-653.
 16. Gaisberger H, Legay S, Andre C, Loo J, Azimov R, Aaliev S, et al. Diversity under threat: Connecting genetic diversity and threat mapping to set conservation priorities for *Juglans regia* L. populations in Central Asia. *Front Ecol Evol.* 2020;8:171.
 17. Maua JO, Mugatsia Tsingalia H, Cheboiwo J, Odee D. Population structure and regeneration status of woody species in a remnant tropical forest: A case study of South Nandi forest, Kenya. *Glob Ecol Conserv.* 2020;21:e00820.
 18. Ahmed AT, Desougi MA. The Ecology of Some Gum Tree Species and Their Uses at Tozi Natural Forest, Sinnar State, Sudan. *J Nat Resour Environmental Stud.* 2014;6456:29-35.
 19. Deng B, Tammeorg P, Luukkanen O, Helenius J, Starr M. Effects of *Acacia seyal* and biochar on soil properties and sorghum yield in agroforestry systems in South Sudan. *Agrofor Syst.* 2017;91:137-148.
 20. Beche D, Gebeyehu G, Feyisa K. Indigenous utilization and management of useful plants in and around Awash National Park, Ethiopia. *J Plant Biol Soil Heal.* 2016;3(1):12.
 21. Louis H, Linus MN, Israt A, Innocent J, Amos PI, Magu TO. Antimicrobial Activity of Stem, Leave and Root Plant Extract of *Sclerocarya birrea* and *Sterculia setigera* against Some Selected Microorganisms. *World Sci News.* 2018;92(2):309-326.
 22. Dampney FG, de la Riva EG, Birkhofer K. Trade-Offs and Synergies Between Food and Fodder Production and Other Ecosystem Services in an Actively Restored Forest, Natural Forest and an Agroforestry System in Ghana. *Front For Glob Chang.* 2021;4:630959.
 23. Heym M, Uhl E, Moshhammer R, Dieler J, Stimm K, Pretzsch H. Utilising forest inventory data for biodiversity assessment. *Ecol Indic.* 2021;121:107196.
 24. Ghanbari S, Sefidi K, Kern CC, Álvarez-Álvarez P. Population structure and regeneration status of woody plants in relation to the human Interventions, Arasbaran Biosphere Reserve, Iran. *Forests.* 2021;12(2):191.
 25. Benítez-Badillo G, Lascunain-Rangel M, Álvarez-Palacios JL, Gómez-Díaz JA, Avendaño-Reyes S, Dávalos-Sotelo R, et al. Influence of Land-Use Changes (1993 and 2013) in the Distribution of Wild Edible Fruits From Veracruz (Mexico). *Trop Conserv Sci.* 2018;11:1940082918758662.
 26. Bentsi-Enchill F, Dampney FG, Pappoe AN, Ekumah B, Akotoye HK. Impact of anthropogenic disturbance on tree species diversity, vegetation structure and carbon storage potential in an upland evergreen forest of Ghana, West Africa. *Trees For People.* 2022;8:100238.
 27. Kimaro J, Lulandala L. Human influences on tree diversity and composition of a coastal forest ecosystem: The case of Ngumburuni forest reserve, Rufiji, Tanzania. *Int J For Res.* 2013;2013:1-7.
 28. Ball L, Tzanopoulos J. Livestock browsing affects the species composition and structure of cloud forest in the Dhofar Mountains of Oman. *Appl Veg Sci.* 2020;23(3):363-376.
 29. Lacerda AE, Kellermann B. What is the Long-Term Effect of Bamboo Dominance on Adult Trees in the *Araucaria* Forest? A Comparative Analysis between Two Successional Stages in Southern Brazil. *Diversity.* 2019;11(9):165.
 30. John E, Bunting P, Hardy A, Roberts O, Giliba R, Silayo DS. Modelling the impact of climate change on Tanzanian forests. *Divers Distrib.* 2020;26(12):1663-1686.
 31. Wang X, Yan J, Zhang X, Zhang S, Chen Y. Organic manure input improves soil water and nutrients use for sustainable maize (*Zea mays*. L) productivity on the Loess Plateau. *PLoS One.* 2020;15(8):e0238042.
 32. Marone D, Poirier V, Coyea M, Olivier A, Munson AD. Carbon storage in agroforestry systems in the semi-arid zone of Niayes, Senegal. *Agrofor Syst.* 2017;91:941-954.
 33. Ouedraogo S, Bonde L, Ouedraogo O, Ouedraogo A, Thiombiano A, Boussim IJ. To what extent do tree size, climate and land use influence the fruit production of *Balanites aegyptiaca* (L) Delile in tropical areas (Burkina Faso)? *Int J Fruit Sci.* 2020;20(3):282-299.
 34. Endale Y, Derero A, Argaw M, Muthuri C. Farmland tree species diversity and spatial distribution pattern in semi-arid East Shewa, Ethiopia. *For Trees Livelihoods.* 2017;26(3):199-214.
 35. Mohammed EM, Hassan TT, Idris EA, Abdel-Magid TD. Tree population structure, diversity, regeneration status, and potential disturbances in Abu Gadaf natural reserved forest, Sudan. *Environ Challenges.* 2021;5:100366.
 36. Hassan TT. The use of non-timber forest products as potential adaptation measures in the Sudan's forestry sector, University of Bahri. 2019.
 37. Abdelrahim M. Contribution of Non-wood Forest Products in Support of Livelihoods of Rural People Living in the Area South of Blue Nile State, Sudan. *Int J Agric For Fish.* 2015;3:189-194.
 38. Mahgoub AA. Changes in Vegetation Cover and Impacts of Human Population in Dinder National Park, Sudan During the Period from 1972 to 2013. *Sudan Academy of Science (SAS).* 2014.
 39. Al Alzubairi MA, Hamdan MM. Deterioration of Blue Nile forests and its ecological effects in the Gezira State– Sudan. *MOJ Ecol Environ Sci.* 2020;5(1):34-40.
 40. El Mamoun HO, El Zein A. Species Dynamics and Potential Disturbances in El Nour Natural Forest Reserve, Sudan. *J For Prod Ind.* 2012;1(2):10-20.
 41. Elmekki AA. Towards community's involvement in integrated management in Dinder Biosphere Reserve, Sudan. University of Khartoum. 2008:1-187.
 42. Deafalla TH, Dafa-Alla DM, Abbas ME. The importance of non wood forest products for rural livelihoods: The case of South Kordofan State, Sudan. *Sci Policy Polit Mod Agric Syst.* 2014:323-332.
 43. Mohammed EM. Impact of land use practices on the natural forest in Dinder Biosphere Reserve and its surrounding areas – Sudan. University of Bahri. 2019.
 44. Hassan TT. Forest Resource Assessment for Transitional Zone of Dinder Biosphere Reserve (DBR)–Sudan. Forestry Consultancy Final Report, Submitted to the Sudanese National Commission for Education, Sciences and Culture (NATCOMUNESCO) and the Wildlife Conservation General Administration (WCGA), Khartoum, Sudan. 2015.
 45. Hassan TT, Mohammed EM, Magid TD. Exploring the current status of forest stock in the areas bordering Dinder Biosphere Reserve, Sudan. *J For Nat Resour.* 2022;1(1):11-25.
 46. Mohammed EM, Hammed AM, Minnick TJ, Ndakidemi PA, Treydte AC. Livestock browsing threatens the survival of *Balanites aegyptiaca* seedlings and saplings in Dinder Biosphere Reserve, Sudan. *J Sustain For.* 2022;41(10):1046-1063.
 47. Ibrahim E, Osman E, Idris E, Yousif T. Linear and Non-Linear Regression Equations for Estimating the Crown Diameter of Three Sudanese Edible Trees. *J For Prod Ind.* 2015;4(2):44-52.
 48. Mohammed EM, Elhag AM, Ndakidemi PA, Treydte AC. Anthropogenic pressure on tree species diversity, composition, and growth of *Balanites aegyptiaca* in Dinder Biosphere Reserve, Sudan. *Plants.* 2021;10(3):483.
 49. Gebeyehu G, Soromessa T, Bekele T, Teketay D. Species composition, stand structure, and regeneration status of tree species in dry Afromontane forests of Awi Zone, northwestern Ethiopia. *Ecosyst Health Sust.* 2019;5(1):199-215.
 50. Ligate EJ, Wu C, Chen C. Investigation of tropical coastal forest regeneration after farming and livestock grazing exclusion. *J For Res.* 2019;30(5):1873-1884.
 51. Kikoti IA, Mligo C, Kilemo DB. The impact of grazing on plant natural regeneration in northern slopes of Mount Kilimanjaro, Tanzania. *Open J Ecol.* 2015;5(6):266.
 52. Hasoba AM, Siddig AA, Yagoub YE. Exploring tree diversity and stand structure of savanna woodlands in southeastern Sudan. *J Arid Land.* 2020;12:609-617.

53. Mekuria W, Aynekulu E. Exclosure land management for restoration of the soils in degraded communal grazing lands in northern Ethiopia. *Land Degrad Dev.* 2013;24(6):528-538.
54. Scholten T, Goebes P, Kühn P, Seitz S, Assmann T, Bauhus J, et al. On the combined effect of soil fertility and topography on tree growth in subtropical forest ecosystems—a study from SE China. *J Plant Ecol.* 2017;10(1):111-127.
55. Suleiman MS, Wasonga VO, Mbau JS, Suleiman A, Elhadi YA. Non-timber forest products and their contribution to households income around Falgore Game Reserve in Kano, Nigeria. *Ecol Process.* 2017;6:1-14.
56. Adam YO, Pretzsch J, Pettenella D. Contribution of Non-Timber Forest Products livelihood strategies to rural development in drylands of Sudan: Potentials and failures. *Agric Syst.* 2013;117:90-97.
57. Ereso T. The role of *Faidherbia albida* tree species in parkland agroforestry and its management in Ethiopia. *J Hortic For.* 2019;11(3):42-47.
58. Gebru BM, Wang SW, Kim SJ, Lee WK. Socio-ecological niche and factors affecting agroforestry practice adoption in different agroecologies of southern Tigray, Ethiopia. *Sustainability.* 2019;11(13):3729.
59. Khamis G, Saleh AM, Habeeb TH, Hozzein WN, Wadaan MA, Papenbrock J, et al. Provenance effect on bioactive phytochemicals and nutritional and health benefits of the desert date *Balanites aegyptiaca*. *J Food Biochem.* 2020;44(6):e13229.
60. Piabuo SM, Minang PA, Tieguhong CJ, Foundjem-Tita D, Nghobuoche F. Illegal logging, governance effectiveness and carbon dioxide emission in the timber-producing countries of Congo Basin and Asia. *Environ Dev Sustain.* 2021;23:14176-14196.
61. Bayih W, Yihune M. Population status, feeding ecology and activity pattern of common bushbuck (*Tragelaphus scriptus decula*) in Sekele Mariam Forest, West Gojjam, Ethiopia. *J Ecol Nat Environ.* 2018;10(5):69-79.
62. Idrissa B, Soumana I, Issiaka Y, Karimou AJ, Mahamane A, Mahamane S, et al. Trend and structure of populations of *balanites aegyptiaca* in parkland agroforests in Western Niger. *Annu Res Rev Biol.* 2018;1-12.
63. Ji Y, Ranjan R, Truong C. Determinants of illegal logging in Indonesia: An empirical analysis for the period 1996-2010. *J Sustain For.* 2018;37(2):197-220.
64. Chaudhary A, Burivalova Z, Koh LP, Hellweg S. Impact of forest management on species richness: global meta-analysis and economic trade-offs. *Sci Rep.* 2016;6(1):23954.
65. Adekunle VA, Olagoke AO. The impacts of timber harvesting on residual trees and seedlings in a tropical rain forest ecosystem, southwestern Nigeria. *Int J Biodivers Sci Ecosyst Serv Manag.* 2010;6(3-4):131-138.
66. Fakhry AM, Aljedaani GS. Impact of disturbance on species diversity and composition of *Cyperus conglomeratus* plant community in southern Jeddah, Saudi Arabia. *J King Saud Univ Sci.* 2020;32(1):600-605.
67. Aleza K, Villamor GB, Nyarko BK, Wala K, Akpagana K. Shea (*Vitellaria paradoxa* Gaertn CF) fruit yield assessment and management by farm households in the Atacora district of Benin. *PloS One.* 2018;13(1):e0190234.
68. Cheng SH, Ahlroth S, Onder S, Shyamsundar P, Garside R, Kristjanson P, et al. What is the evidence for the contribution of forests to poverty alleviation? A systematic map protocol. *Environ Evid.* 2017;6(1):1-11.
69. Le Houérou HN. Utilization of fodder trees and shrubs in the arid and semiarid zones of West Asia and North Africa. *Arid Soil Res Rehabil.* 2000;14(2):101-135.
70. Belant JL, Bled F, Mkasanga JJ, Wilton CM, Mwampeta SB, Beyer DE, et al. Track surveys do not provide accurate or precise lion density estimates in serengeti. *Glob Ecol Conserv.* 2019;19:e00651.
71. Amahowe IO, Gaoe OG, Natta AK, Piponiot C, Zobi IC, Hérault B. Functional traits partially mediate the effects of chronic anthropogenic disturbance on the growth of a tropical tree. *AoB Plants.* 2018;10(3):ply036.
72. Ahenkan A, Boon E. Improving nutrition and health through non-timber forest products in Ghana. *J Health Popul Nutr.* 2011;29(2):141-148.
73. Bondé L, Ouédraogo O, Ouédraogo I, Thiombiano A, Boussim JI. Variability and estimating in fruiting of shea tree (*Vitellaria paradoxa* CF Gaertn) associated to climatic conditions in West Africa: implications for sustainable management and development. *Plant Prod Sci.* 2019;22(2):143-158.
74. Ali OB, Ibrahim EM, Abdelmagid TD. Detection of tree species dynamic changes on Savannah ecology through diameter at breast height, Case study Heglieg area, Sudan. *J Nat Resour Environ Stud.* 2015;3(1):24-30.
75. Heshmati M, Gheitury M, Hosseini M. Effects of runoff harvesting through semi-circular bund on some soil characteristics. *Glob J Environ Sci Manag.* 2018;4(2):207-216.
76. Mao P, Han G, Wang G, Yu J, Shao H. Effects of age and stand density of mother trees on early *Pinus thunbergii* seedling establishment in the Coastal Zone, China. *Sci World J.* 2014;2014:1-9.
77. Ritsche J, Katzensteiner K, Acácio V. Tree regeneration patterns in cork oak landscapes of Southern Portugal: The importance of land cover type, stand characteristics and site conditions. *For Ecol Manag.* 2021;486:118970.
78. Singh S, Malik ZA, Sharma CM. Tree species richness, diversity, and regeneration status in different oak (*Quercus* spp.) dominated forests of Garhwal Himalaya, India. *J Asia Pac Biodivers.* 2016;9(3):293-300.
79. Jayakumar R, Nair KK. Species diversity and tree regeneration patterns in tropical forests of the Western Ghats, India. *Int Sch Res Notices.* 2013;2013.
80. Crump L, Mauti S, Traoré A, Shaw A, Hattendorf J, Zinsstag J. The contribution of livestock to urban resilience: the case of Bamako, Mali. *Trop Anim Health Prod.* 2019;51:7-16.
81. Blackham GV, Thomas A, Webb EL, Corlett RT. Seed rain into a degraded tropical peatland in Central Kalimantan, Indonesia. *Biol Conserv.* 2013;167:215-223.
82. Ewunetu A, Simane B, Teferi E, Zaitchik BF. Land cover change in the blue Nile river headwaters: farmers' perceptions, pressures, and satellite-based mapping. *Land.* 2021;10(1):68.
83. Dayamba SD, Tigabu M, Sawadogo L, Oden PC. Seed germination of herbaceous and woody species of the Sudanian savanna-woodland in response to heat shock and smoke. *For Ecol Manag.* 2008;256(3):462-470.
84. Liu J, Bai X, Yin Y, Wang W, Li Z, Ma P. Spatial patterns and associations of tree species at different developmental stages in a montane secondary temperate forest of northeastern China. *Peer J.* 2021;9:e11517.
85. Cantarello E, Lovegrove A, Orozumbekov A, Birch J, Brouwers N, Newton AC. Human impacts on forest biodiversity in protected walnut-fruit forests in Kyrgyzstan. *J Sustain For.* 2014;33(5):454-481.
86. Neelo J, Teketay D, Kashe K, Masamba W. Stand structure, diversity and regeneration status of woody species in open and enclosed dry woodland sites around Molapo farming areas of the Okavango Delta, Northeastern Botswana. *Open J For.* 2015;5(4):313-328.
87. Wang S, Fan J, Li Y, Huang L. Effects of grazing exclusion on biomass growth and species diversity among various grassland types of the Tibetan Plateau. *Sustainability.* 2019;11(6):1705.
88. de la Peña-Domene M, Howe HF, Cruz-León E, Jiménez-Rolland R, Lozano-Huerta C, Martínez-Garza C. Seed to seedling transitions in successional habitats across a tropical landscape. *Oikos.* 2017;126(3):410-419.
89. Asigbaase M, Sjogersten S, Lomax BH, Dawoe E. Tree diversity and its ecological importance value in organic and conventional cocoa agroforests in Ghana. *PLoS One.* 2019;14(1):e0210557.
90. Amiri F, Ariapour A, Fadai S. Effects of livestock grazing on vegetation composition and soil moisture properties in grazed and

- non-grazed range site. *J Biol Sci.* 2008;8(8):1289-1297.
91. Reuter KE, Sewall BJ, Di Minin E. Drivers of present and lifetime natural resource use in a tropical biodiversity hotspot. *Anim Conserv.* 2018;21(2):127-136.
92. Arosa ML, Bastos R, Cabral JA, Freitas H, Costa SR, Santos M. Long-term sustainability of cork oak agro-forests in the Iberian Peninsula: a model-based approach aimed at supporting the best management options for the montado conservation. *Ecol Modell.* 2017;343:68-79.
93. Aubad J, Aragón P, Olalla-Tárraga MÁ, Rodríguez MÁ. Illegal logging, landscape structure and the variation of tree species richness across North Andean Forest remnants. *For Ecol Manag.* 2008;255(5-6):1892-1899.
94. Kutnar L, Nagel TA, Kermavnar J. Effects of disturbance on understory vegetation across Slovenian forest ecosystems. *Forests.* 2019;10(11):1048.
95. Tetemke BA, Birhane E, Rannestad MM, Eid T. Allometric models for predicting aboveground biomass of trees in the dry afro-montane forests of Northern Ethiopia. *Forests.* 2019;10(12):1114.
96. Wang Y, Yu J, Xiao L, Zhong Z, Wang Q, Wang W. Dominant species abundance, vertical structure and plant diversity response to nature forest protection in Northeastern China: Conservation effects and implications. *Forests.* 2020;11(3):295.