

EDITORIAL



Integrating health education on diabetes prevention for young people in higher institutions

Foluke Stella Bosun-Arije

Department of Nursing, University of Doha for Science and Technology, Doha, Qatar

ARTICLE HISTORY

Received 02 June 2026; Accepted 23 June 2026; Published 1 July 2026

Diabetes mellitus (DM) is a chronic metabolic condition characterized by elevated blood glucose levels due to the body's inability to produce sufficient insulin or effectively utilize the insulin produced [1,2]. The prevalence and incidence of DM are increasing globally. In 2024, approximately 588.7 million individuals worldwide were living with DM, and projections indicate that this number could rise to 852.5 million individuals aged 20 to 79 years by 2050 if current trends continue [3]. A survey conducted in South Africa revealed a prevalence of pre-diabetes and DM at 67% and 22%, respectively [4]. Among those who had never been tested for DM prior to the survey, 10% of females and 6% of males were found to have pre-diabetes, with 67% of both males and females exhibiting pre-diabetic conditions [4]. The International Diabetes Federation emphasizes that pre-diabetes predisposes young individuals to a high risk of developing Type 2 Diabetes (T2D), underscoring DM as one of the fastest-growing global health emergencies.

Accumulating evidence indicates that among the three main types of DM, T2D is the most preventable yet the most prevalent, accounting for 90% of all global DM cases. Notably, three out of four adults living with T2D reside in low- and middle-income countries (LMICs), with approximately 59% of DM cases remaining undiagnosed despite the availability of affordable and proven preventive measures. The onset of T2D is often asymptomatic, making it difficult to determine, and unmanaged T2D can lead to severe complications [5,6].

Although specific epidemiological data on the number of individuals living with T2D in LMICs are lacking, the burden of T2D has overwhelmed many healthcare systems in these regions and poses an increased financial burden on high-income countries [7]. In the United Kingdom, the National Health Service (NHS) allocates approximately £10 billion annually to DM and its related complications, equivalent to 10% of the national budget [8]. Sedentary lifestyles, poor eating habits, and low physical activity levels are closely linked to T2D [9]. Additionally, high consumption of sweetened beverages and unhealthy foods are contributing factor to obesity, a precursor to T2D [10].

DM education remains a cornerstone for the prevention and management of pre-diabetes and T2D. Educational initiatives promote the prevention of obesity and its contributing factors, making education a pivotal element in T2D management and pre-diabetes mitigation. In India, the

Prediabetes Education Programme (PEP) significantly improved the knowledge and attitudes of individuals with pre-diabetes towards lifestyle modification [11]. Education encourages lifestyle changes, healthy eating, regular physical activity to halt pre-diabetes progression, and smoking cessation to improve glycaemic profiles [12-15].

Reducing alcohol consumption aids in maintaining a healthy body weight [16]. Diabetes remission was observed in 61% of participants in an intervention group who received education and achieved an annual body weight reduction of about 12kg [17]. However, Baker et al. caution that the content, modality, setting, and providers of DM education should ensure the delivery of high-quality and comprehensive DM education [18]. Therefore, when educating young individuals in higher education, DM educators should collaborate with academics to deliver effective DM education.

Young individuals often juggle school and work commitments and may seek health information on social media. While accurate health information can be obtained via social media, there is a risk of encountering misleading and non-evidence-based information. Research conducted across Europe, the USA, Pakistan, Canada, and India by Horigan et al. suggests that the busy lifestyles of young individuals may explain the non-attendance of 2,260 young patients with diabetes at scheduled structured DM educational sessions. Zowgar et al. conducted a DM knowledge test for 942 patients in Saudi Arabia, including 209 university-level young adults (28.1%), all of whom demonstrated a lack of DM literacy [19-21].

Structured DM education within routine practice has shown clinically significant benefits for young individuals in Australia. In Nigeria, Bosun-Arije et al. identified knowledge and information deficiencies as factors contributing to the high prevalence and incidence of T2D [2]. The high prevalence of overweight and obesity is a significant public health concern that exceeds geographical boundaries within Africa, necessitating innovative weight-management education and interventions for young individuals [22-24].

T2D can be silent and progressive. Epidemiological analyses suggest that identifying and educating populations and subgroups at risk of T2D and its complications is an innovative approach to curbing the condition [25]. If left unchecked, T2D can lead to a range of microvascular and macrovascular

complications, posing a significant health challenge to young individuals, families, healthcare facilities, and societies. If ignored, up to 40% of individuals diagnosed with T2D may develop chronic kidney disease (CKD), placing them at higher risk of cardiovascular disease (CVD), kidney failure, and fatality [7].

This editorial advocates for the importance of diabetes education in higher education to enhance DM literacy and awareness among young individuals. These efforts can provide a pragmatic approach to T2D prevention, aligning with the IDF strategic agenda, which aims to keep T2D in the public spotlight and promote coordinated and concerted actions to address the impacts of T2D and other types of DM in the 21st century [7,26].

References

- Global Burden of Disease Collaborative Network. Global Burden of Disease Study 2019. Results. Institute for Health Metrics and Evaluation. 2020. Available at <https://vizhub.healthdata.org/gbd-results/>.
- World Health Organisation 2023. Diabetes and Key Facts. Available at: <https://www.who.int/news-room/fact-sheets/detail/diabetes>.
- International Diabetes Federation (2025). IDF Atlas 11th edition. Available at: <https://diabetesatlas.org/resources/idf-diabetes-atlas-2025/>.
- Grundlingh N, Zewotir TT, Roberts DJ, Manda S. Assessment of prevalence and risk factors of diabetes and pre-diabetes in South Africa. *J Health Popul Nutr*. 2022;41(1):7. <https://doi.org/10.1186/s41043-022-00281-2>
- Gregg EW, Li Y, Wang J, Rios Burrows N, Ali MK, Rolka D, et al. Changes in diabetes-related complications in the United States, 1990–2010. *N Engl J Med*. 2014;370(16):1514–1523. <https://doi.org/10.1056/NEJMoa1310799>
- King P, Peacock I, Donnelly R. The UK prospective diabetes study (UKPDS): clinical and therapeutic implications for type 2 diabetes. *Br J Clin Pharmacol*. 1999;48(5):643. <https://doi.org/10.1046/j.1365-2125.1999.00092.x>
- International Diabetes Federation. IDF Diabetes Atlas Report 2023. Available at: <https://diabetesatlas.org/atlas-reports/?report-year=2023>.
- Whicher CA, O'Neill S, Holt RG. Diabetes in the UK: 2019. *Diabetic Medicine*. 2020;37(2):242–247. <https://doi.org/10.1111/dme.14225>
- Lascar N, Brown J, Pattison H, Barnett AH, Bailey CJ, Bellary S. Type 2 diabetes in adolescents and young adults. *Lancet Gastroenterol Hepatol*. 2018;6(1):69–80.
- Bellou V, Belbasis L, Tzoulaki I, Evangelou E. Risk factors for type 2 diabetes mellitus: an exposure-wide umbrella review of meta-analyses. *PloS one*. 2018;13(3):e0194127. <https://doi.org/10.1371/journal.pone.0194127>
- Hyder KM, Mohan J, Varma V, Ponnusankar S, Raja D. Impact of prediabetes education program on Knowledge, attitude and practice among prediabetic population of south India. *Prev Med Rep*. 2021;23:101395. <https://doi.org/10.1016/j.pmedr.2021.101395>
- McFarlane SI, Seixas AA, Agyemang C, Jean-Louis G. Effects of lifestyle modifications and dietary habits on prevention of diabetes and cardiovascular disease. *J Diabet Res*. 2018;2018:2341230. <https://doi.org/10.1155/2018/2341230>
- American Diabetes Association. 4. Lifestyle management. *Diabetes care*. 2017;40(Supplement_1):S33–S43. <https://doi.org/10.2337/dc17-S007>
- Liu G, Hu Y, Zong G, Pan A, Manson JE, Rexrode KM, et al. Smoking cessation and weight change in relation to cardiovascular disease incidence and mortality in people with type 2 diabetes: a population-based cohort study. *Lancet Diabetes Endocrinol*. 2020;8(2):125–133.
- Durlach V, Vergès B, Al-Salameh A, Bahougne T, Benzerouk F, Berlin I, et al. Smoking and diabetes interplay: a comprehensive review and joint statement. *Diabetes & metabolism*. 2022;48(6):101370. <https://doi.org/10.1016/j.diabet.2022.101370>
- Mayl JJ, German CA, Bertoni AG, Upadhyaya B, Bhavne PD, Yeboah J, et al. Association of alcohol intake with hypertension in type 2 diabetes mellitus: the ACCORD trial. *J Am Heart Assoc*. 2020;9(18):e017334. <https://doi.org/10.1161/JAHA.120.017334>
- Taheri S, Zaghloul H, Chagoury O, Elhadad S, Ahmed SH, El Khatib N, et al. Effect of intensive lifestyle intervention on bodyweight and glycaemia in early type 2 diabetes (DIADEM-I): an open-label, parallel-group, randomised controlled trial. *Lancet Diabetes Endocrinol*. 2020;8(6):477–489.
- Baker KM, Nassar CM, Baral N, Magee ME. The current diabetes education experience: findings of a cross-sectional survey of adults with type 2 diabetes. *Patient Educ Couns*. 2023;108:107615. <https://doi.org/10.1016/j.pec.2022.107615>
- Fergie G, Hilton S, Hunt K. Young adults' experiences of seeking online information about diabetes and mental health in the age of social media. *Health Expectations*. 2016;19(6):1324–1335. <https://doi.org/10.1111/hex.12430>
- Horigan G, Davies M, Findlay-White F, Chaney D, Coates V. Reasons why patients referred to diabetes education programmes choose not to attend: a systematic review. *Diabetic Medicine*. 2017;34(1):14–26. <https://doi.org/10.1111/dme.13120>
- Zowgar AM, Siddiqui MI, Alattas KM. Level of diabetes knowledge among adult patients with diabetes using diabetes knowledge test. *Saudi medical journal*. 2018;39(2):161. <https://doi.org/10.15537/smj.2017.2.21343>
- Speight J, Holmes-Truscott E, Harvey DM, Hendrickx C, Hagger VL, Harris SE, et al. Structured type 1 diabetes education delivered in routine care in Australia reduces diabetes-related emergencies and severe diabetes-related distress: The OzDAFNE program. *Diabetes Res Clin Pract*. 2016;112:65–72. <https://doi.org/10.1016/j.diabres.2015.11.002>
- Bosun-Arije FS, Ling J, Graham Y, Hayes C. Organisational factors influencing non-pharmacological management of type 2 diabetes mellitus (T2DM) in public hospitals across Lagos, Nigeria: A qualitative study of nurses' perspectives. *Diabetes Res Clin Pract*. 2020;166:108288. <https://doi.org/10.1016/j.diabres.2015.11.002>
- Ekpor E, Akyirem S, Adade Duodu P. Prevalence and associated factors of overweight and obesity among persons with type 2 diabetes in Africa: a systematic review and meta-analysis. *Ann Med*. 2023;55(1):696–713. <https://doi.org/10.1080/07853890.2023.2182909>
- Chan JC, Lim LL, Wareham NJ, Shaw JE, Orchard TJ, Zhang P, et al. The Lancet Commission on diabetes: using data to transform diabetes care and patient lives. *The Lancet*. 2020;396(10267):2019–2082.
- Diabetes UK. The Future of Diabetes. 2019. Available at <https://www.diabetes.org.uk/support-us/campaign/other-campaigns/the-future-of-diabetes>.